CHAPTER N0.

03

# Igniting Recovery: The Power of Heat Therapy in Physical Therapy

MEERAB WASEEM[1], WAFA MAJEED[2]*, UJALA MALIK[3], NOOR UN NISA[4], RABEET ZAIMA KHAN[4], ALEENA SHABBIR[4], HANIYA[4]

[1]Department of Epidemiology and Public Health, University of Agriculture, Faisalabad, Pakistan
[2]Department of Pharmacy, University of Agriculture, Faisalabad, Pakistan
[3]Department of Global Public Health, Manchester Metropolitan University, Manchester, England
[4]Department of Rehabilitation Sciences, University of Faisalabad, Faisalabad, Pakistan
*Corresponding author: wafa.majeed@uaf.edu.pk

**Summary**: Thermotherapy, also known as heat therapy, remains a key element in physical rehabilitation due to its broad physiological and functional effects. The chapter is a review of the mechanisms of therapeutic heating, clinical uses, and research evidence that support therapeutic heating. Heat application enhances muscle tension, improves collagen elasticity, stimulates tissue metabolism, and enhances vasodilation, an ideal setting to create a perfect recovery setting. The two superficial methods, such as the use of hot packs, paraffin applications, and fluid therapy and the deep methods, such as ultrasound and shortwave diathermy, are all addressed in terms of their mechanism, clinical indications, and safety measures. Their applicability in the treatment of arthritis, chronic musculoskeletal pain, post-operative stiffness, and as a preparatory device to stretching or exercise is effective. Besides, the chapter also highlights the significance of patient education, contraindication awareness, and heat therapy in combination with exercise, manual techniques, and multidisciplinary care. Although it is widely used, additional research is justified to make treatment guidelines more consistent and clearer regarding the long-term outcomes. Altogether, thermotherapy is a safe, cost-effective and versatile intervention, which improves the quality of life and functions.
**Keywords:** Heat therapy, Thermotherapy, Rehabilitation, Musculoskeletal disorders, Physical therapy, Pain management

## INTRODUCTION

The role of physical therapy in rehabilitation medicine is becoming established as not just a transient measure to relieve any short-term symptoms, but as an approach to achieving more lasting outcomes, like the reintegration of tissue integrity, neuromuscular pathway retraining, and involvement in everyday activities. In this realization, the World Health Organization has seen rehabilitation as a significant public health intervention in enhancing functionality and reducing disability in a wide range of health impairments (World Health Organization, 2024). Among them, musculoskeletal (MSK) disorders that include muscle, joint, bone, and tendon disorders are of importance. They are responsible to a large extent for disability and pain in the world, reducing the quality of life of the impacted people. Available estimates indicate that more than 1.7 billion individuals have MSK disorders, which are expected to increase as the population and the elderly population age (Blyth et al., 2019; World Health Organization, 2024b).

MSK problems are the most common reasons behind up to one-third of general practice (GP) visits, and it is also a commonly used excuse by people to visit a community pharmacy and seek advice. Because of numerous advantages, such as the reduction of pain, the reduction of stiffness and spasms, and the increased flexibility and range of motion, the heat treatment (HT) is an effective alternative method of MSK pain treatment. All said and done, such effects help to promptly resume normal functioning and participation in daily activities and work (Clijsen et al., 2022; Lubrano et al., 2023). Some of the methods through which HT can be used are heat lamps, diathermy, ultrasounds, as well as superficial heat pads or wraps. It is worth mentioning that an HT, in the shape of low-level heat wraps, may be a good alternative used at home and a cost-effective way of providing treatment both in inpatient and outpatient care (Rossi, 2024).

Most ancient civilizations employed bathing, hot springs and topical use of heat and cold in treatment and symptom management; these methods eventually evolved to the balneotherapy and hydrotherapy methods that remain a component of modern rehabilitation and wellness treatments. Among the oldest forms of therapeutic interventions in medicine are the so-called thermotherapy (use of heat) and cryotherapy (use of cold), both of which are known as thermal therapies. The enduring nature of thermal practice can be used to explain its historical availability in modern practicing clinics, in which the use of modern equipment and evidence-based practices has substituted folk practices but still has the same physiological goal (Gianfaldoni et al., 2017).

In contemporary rehabilitation, clinicians use thermal interventions based on the purpose of treatment and the phase of tissue recovery. Cryotherapy is usually recommended in the acute period of the injury because it has vasoconstrictive and anti-inflammatory effect, in the subacute and chronic periods, the heat modalities are more often used to stimulate

How to Cite: Waseem M, W Majeed, U Malik, NU Nisa, RZ Khan, A Shabbir, Haniya, 2025. Igniting recovery: the power of heat therapy in physical therapy. In: Advances in Animal and Life Sciences (Sindhu ZuD, M Imran, MA Zafar, S Yousaf, Eds). FahumSci, Lahore, Pakistan, pp: 16-22. ISBN: 978-627-7745-12-7. https://doi.org/10.61748/AALS.2025/03

circulation, decrease joint stiffness, and improve tissue elasticity. Increasing clinical practice and mechanistic research evidence has given increased knowledge regarding the optimal uses of thermotherapy, such as incorporating heat as a pre-stretching or pre-manual therapy use or in multimodal pain management systems of patients with chronic musculoskeletal conditions (Brunt and Minson, 2021).

The chapter focuses on thermotherapy: the biophysical principles, the main mechanisms of action, clinical indications, the most common delivery techniques, and safety issues. The impartiality is to provide a pragmatic synthesis, which connects patient-centered decision-making automatous knowledge so that heat can be used in a manner that will predictively lead to recovery and functional restoration.

## HEAT THERAPY: PRINCIPLES AND AIMS

Thermotherapy can be defined as using a controlled rise in the temperature of tissues in order to attain therapeutic effects. The clinical goal is to elevate the tissue temperature locally to a range that promotes beneficial physiological responses that include increased blood flow, neuromuscular alterations, increased connective tissue flexibility, and prevention of thermal injury. Surface-heating modalities (e.g., diathermy or ultrasound) are mainly used to influence the skin and subcutaneous layers; conversion-based modalities (e.g., diathermy or ultrasound) are used in the case of reaching deeper tissues. According to the recent reviews, the safe temperature ranges were defined as well as the physiological rationale behind using superficial and deep heating methods (Freiwald et al., 2021).

Heat applications have a number of related functions in clinical practice. First of all, it reduces muscular guarding and changes the nociceptive transmission, which causes analgesic effects and allows engaging actively in treatment. Also, heating of tissues reduces tissue stiffness by altering the viscoelastic properties of collagen-rich structures, thus permitting further stretching or movement to provide more pronounced and long-lasting increases in joint range of motion. Simultaneously, localized circulation and metabolic enhancement facilitate reparative processes, such as fibroblast growth and angiogenesis, especially in subacute and chronic traumas or tendinopathy. The warmth is also related to the comfort of the patient, increasing the adherence to rehabilitation programs, decreasing anxiety and promoting mobility in a patient. Collectively, these outcomes explain the prevalence of the use of thermotherapy by clinicians as a preparatory or complementary intervention and not as a treatment (Charkoudian, 2010).

### Physiological Impact of Heat Therapy

The heat clinical paraphernalia has a foundation on physiologically reproducible reactions functioning at the vascular, connective-tissue, neuromuscular and metabolic systems. Local warming can trigger local axon reflexes and nitric-oxide dependent vasodilatory responses, and local thermoreceptors and reflex, which raise blood flow to the skin and tissue to supply oxygen and nutrients to anaerobic tissue and reduce metabolic by-products, which are mediated by local axon reflexes and nitric-oxide dependent vasodilatory processes. Clinical and experimental studies utilizing Doppler and near-infrared methods have shown that deeper tissue perfusion can be enhanced with deep-heating methods that have been carefully selected and confirm that cutaneous perfusion can increase with localized heating (Rutkove, 2001).

Mechanical comportment of collagenous tissues is also altered with heating. The viscoelasticity of collagen is higher at high tissue temperature, which implies that the substance becomes less firm. In a combination of heat and controlled stretching, a measurable and clinically important effect is observed in the improvement in joint range of motion and decrease in stiffness. This interaction supports the use of common clinical procedures that involve heat in advance of mobilization or stretching to treat adhesive capsulitis, persistent contractures, or post-immobilization stiffness. (Hill et al., 2024).

There is a variation of analgesic effects of heat. It is estimated that the combined action of heat, eliminating muscle spindle-mediated guarding by diminishing gamma efferent drive and non-noxious afferent input (e.g., thermal signals), inhibiting nociceptive transmission at the spinal and supraspinal levels is fast subjective pain relief, enhanced activity tolerance and enhanced involvement in active rehabilitation, in accordance with classical and modern models of pain modulation. Such a neuromodulator action, consisting of local ischemia reductions by vasodilation, has aided in explaining the role of heat in chronic low back pain, osteoarthritis and other chronic musculoskeletal pain syndromes (Ventriglia et al., 2023).

At the cellular level, heat facilitates matrix remodeling and fibroblast hyperplasia in subacute and chronic repair by an amplified metabolic rate and enzymatic activity in a temperature-sensitive fashion. The timing is also important in that the sooner the heat is introduced into an inflammatory cascade, the worse bleeding or edema is likely to grow, even though the slightest metabolic acceleration can lead to healing at later stages. The reason clinicians tend to implement heat preceding low-to-moderate intensity exercise over maximal exercise is that the neural conduction velocity and muscle performance are sensitive to temperatures. Warming tends to elevate the conduction velocity and can temporarily change the strength and fatigue properties (Petrofsky et al., 2015).

Although deep modalities of heating such as shortwave diathermy or therapeutic ultrasound enhance perfusion in deeper tissues, such as tendons and joint capsules, recent studies that use near-infrared spectroscopy (NIRS) and Doppler ultrasound demonstrate that superficial heat (such as hot packs) can enhance skin blood flow up to 400 -450 percent of the control condition as shown in Table 1 (Petrofsky et al., 2015).

In conclusion, thermotherapy can provoke interstitial fluid flows and the lymphatic circulation. Although the literature is not uniformly conclusive, there are systematic reviews that supply conditional evidence that appropriately controlled

**Table 1.** Physiological effects of heat therapy: mechanism and clinical implications

| Physiological Effect | Mechanism | Clinical Implication | References |
|---|---|---|---|
| Vasodilation and ↑ Blood Flow | Thermoreceptor activation; nitric oxide release | Improved oxygenation, waste removal, and faster healing | Charkoudian, 2010 |
| Tissue Extensibility | Reduced collagen viscosity; cross-link breakdown | Improved ROM; reduced stiffness in joints/tendons | Petrofsky et al., 2015 |
| Pain Relief | Gate control, pain threshold, spasm | Symptom relief in arthritis, back pain, and fibromyalgia | Nadler et al., 2002 |
| Metabolic Rate | 10–13% rise per 1°C | Accelerated repair; fibroblast activity, angiogenesis | Lehmann et al., 1974 |
| Nerve Conduction Velocity | 1.5–2.0 m/s rise per 1°C | Reduced spasm, improved neuromuscular coordination | Rutkove, 2001 |
| Lymphatic and Fluid Velocity | Lowered viscosity; improved lymphatic drainage | Edema resolution; enhanced recovery | Brunt and Minson, 2021 |
| Muscle Strength (variable) | Altered excitability; reduced fatigue | Warm-up: improved endurance in submaximal tasks | Petrofsky et al., 2015 |

heating can help in chronic lymphedema (for example, as an adjunct to decongestive methods), although further, higher-quality trials are required to determine optimal dosing and safety margins (Hill et al., 2024) (Table 1).

## APPLICATION TECHNIQUES AND MODALITIES

The therapeutic efficacy of heat therapy depends on the physiological responses to heat and the route of administration of heat. The modality used influences the depth of penetration, duration of therapeutic action, speed of heat transfer, comfort to the patient and clinical outcome. Physical therapists should thus be careful in the selection of their techniques, with a consideration of the target organ, the progression of the condition, and other patient-related factors such as comorbidity or skin sensitivity (Alghadir et al., 2019; Vrindten et al., 2025).

### Heat Transfer Mechanisms

The therapeutic use of heat in rehabilitation requires several physical means of heat transfer, with each providing a different clinical advantage. Conduction represents the most familiar approach, where the heat is transported directly by a warmer surface to the cooler tissues via physical contact. Examples of this technique include heating pads, hot packs, and paraffin baths and it works best on superficial tissue with a depth of about one centimeter. By dissimilarity, convection is a process that is used to transfer heat to the body surface by moving particles or fluids across the surface. The modalities, such as fluid therapy and whirlpool therapy, prove this process, particularly the distal extremity and unequal anatomical surfaces. They also facilitate mobilization exercises.

A less commonly mentioned mechanism is evaporation, where the heat is lost as the liquid evaporates at the surface of the skin. Cooling with vapocoolant sprays is mostly associated with evaporation, which can also indirectly support wet-heat applications, in which surface warming is facilitated by humidity. The other type of heat transfer is radiative, which involves the use of electromagnetic waves and does not need physical contact. Conventional ones are far-infrared saunas and infrared lamps; they can only reach the superficial tissues, but they can effectively treat large areas of the surface and gain popularity in sports and wellness facilities (Peng et al., 2020).

Finally, there is a conversion to convert non-thermal energy to heat within the tissues of the body. Such modalities as therapeutic ultrasound, converting acoustic energy, and shortwave diathermy, converting electromagnetic energy, are grounded on this concept. These are especially applicable in the deeper structures like muscles, tendons and joint capsules where the heat penetration may go up to three to five centimeters. By comprehending the combination of these pathways, physicians are able to select the most appropriate modality to benefit a single patient and reduce risk by maximizing therapeutic outcomes (Table 2) (Lehmann et al., 1974; Vrindten et al., 2025).

### Superficial Heat Modalities

Superficial heating methods continue to remain a foundation of clinical and home-based rehabilitation practice. Pragmatic through conduction, heating pads are typically used in mild musculoskeletal pain, and in menstrual pain. They are fast, cheap and convenient, whether electrically or chemically activated. Randomized controlled trials have shown that they are not only helpful in the aspect of comfort, but are competent in relieving chronic low back pain and primary dysmenorrhea and their effects are even comparable to the effects of pharmacological agents such as NSAIDs (Freiwald et al., 2021).

Hot packs, which are conducted, are the most common forms of therapeutic modalities. These are usually made of silica gel that is placed in a fabric, now heated to about 70 to 75 o C and then wrapped in towels and is used to evenly warm large areas of subcutaneous muscle, like the quadriceps or the lower back. There have been reports that hot packs increase tissue pliability with significant analgesic effects, particularly in use with stretching or mobilization procedures (Dogru et al., 2008).

Fluid therapy involves convection heat and it has the benefit that it permits active movement of the treated limb during treatment. The hand, wrist, or foot in this procedure is put in a chamber with hot cellulose, which is then moved around by air currents. Fluid therapy is specifically successful in hand rehabilitation as it integrates all three elements of warmth, mechanical stimulation, and movement. Research has reported an increase in grip strength, functional ability, and pain relief in diseases like osteoarthritis and post-surgical

stiffness, as shown in Table 2 (Oncel et al., 2020; Peng et al., 2020).

Another method of conduction technique is the use of paraffin wax baths, which are mostly used on smaller joints in the hands and feet. In this process, the problem area is repeatedly dipped in a mixture of paraffin and mineral oil kept at approximately 52C and the wax is deposited repeatedly in layers to trap heat within the layers. Arthritis and scleroderma are conditions that are often treated with this intervention to relieve pain, spasms, and swelling. Systematic reviews and clinical studies still confirm its usefulness in the treatment of Osteoarthritis and rheumatoid arthritis as shown in Table 3 (Dilek et al., 2013; Li et al., 2023).

### Deep Heat Modalities

Deep-heating modalities, unlike the superficial heating agents, allow therapeutic energy to penetrate several centimeters beneath the skin, and are thus particularly useful to the structures of tendons, ligaments and joint capsules (Table 1). The ultrasound therapy attains this by transforming acoustic waves into localized thermal energy, where the depth of penetration is based on the frequency, usually around 1 MHz for deep tissue and around 3 MHz for more superficial tissue. It has been extensively documented as having clinical value, particularly in combination with stretching exercises. Draper et al. (1998) established that preoperative ultrasound enhances the shoulder range of motion in people with adhesive capsulitis, and later random studies have validated that ultrasound is effective in the treatment of chronic tendinopathy and postoperative stiffness (Table 2) (Onal Balcı et al., 2018).

Another form of deep heating, shortwave diathermy (SWD), is set into action by the conversion of electromagnetic energy at high frequencies, which is normally at 27.12 MHz, into heat. SWD is also able to penetrate deeper muscle areas and joint structures with large treatment areas, unlike ultrasound. It has been applied effectively in the treatment of chronic low back pain, osteoarthritis, spasm and pain syndromes of the pelvis. Systematic review provides support that it can reduce pain and enhance function and moderate to high levels of support for its use in chronic musculoskeletal conditions (Nascimento et al., 2022).

## CLINICAL APPLICATIONS

Heat therapy (thermotherapy) has a wide variety of uses within the field of rehabilitation, as well as in musculoskeletal medicine. It mostly finds a clinical application in subacute and chronic diseases, after the inflammatory process has passed, and the remodelling of the tissue or pain regulation is the focus point. The action vasodilation, neural modulation, metabolic acceleration, and collagen extensibility apparatuses are consistent with positive patient outcomes, including pain reduction, increased range of motion, and increased meaning (Alghadir et al., 2019; Vrindten et al., 2025).

### Indications for Heat Therapy

Clinical guidelines recommend the use of heat therapy in a variety of clinical scenarios. It is indicated for subacute and chronic musculoskeletal injuries, such as strains and sprains, where it can help alleviate discomfort and improve tissue flexibility. Heat therapy is also beneficial for managing muscle stiffness and spasms associated with neurological or orthopedic conditions. In patients with osteoarthritis or rheumatoid arthritis, particularly during the non-acute phase, heat application can reduce pain and joint stiffness, improving functional mobility. Additionally, heat therapy serves as an adjunct to exercise in the management of fibromyalgia and other widespread chronic pain syndromes. It can also support post-acute tissue healing, especially during the remodeling phase of recovery. Finally, the application of heat before exercise has been shown to enhance muscle extensibility and may reduce the risk of injury, making it a useful component of pre-exercise warm-up routines (Lehmann et al., 1974).

## SPECIFIC CONDITIONS AND INJURIES BENEFITING FROM HEAT

### Chronic Pain Disorders

Thermotherapy consistently improves chronic musculoskeletal pain, which is mostly in the lumbar region, shoulder and neck area. Constant low-level heat wraps have a significant effect in reducing the amount of pain and enhancing functionality (Nadler et al., 2002). Other recent meta-analyses validate the effectiveness of heat therapy in the management of chronic low-back pain, especially when used in conjunction with exercise (Li et al., 2023).

### Muscle Problems

Heat reduces muscle spasm by reducing the excitability of alpha motor neurons, reducing muscle tone, and causing relaxation. It is used in myofascial pain syndrome, cervical tension and delayed-onset muscle soreness (DOMS). Localized heating was found to hasten the recovery of exercise soreness (Petrofsky et al., 2015). Recent research supports the application of moist heat in athletes in order to accelerate recovery and enhance performance metrics (Hill et al., 2024).

### Stiffness and Decreased Range of Motion (ROM)

In other cases, thermodynamics is a supplement to collagen extensibility and includes the treatment of adhesive capsulitis (frozen shoulder) and post-surgical contractures. Dogru et al.

**Table 2.** Comparison of superficial vs. deep heat modalities: mechanism, penetration depth

| Modality | Mechanism | Penetration Depth | Best For | References |
|---|---|---|---|---|
| Heating Pads | Conduction | ≤1 cm | Home-based pain relief, menstrual pain | Nadler et al., 2002 |
| Hot Packs | Conduction | 1 cm | Large superficial areas (e.g., lumbar) | Peng et al., 2020 |
| Fluidotherapy | Convection | Superficial | Hands, feet, active mobilization | Peng et al., 2020 |
| Paraffin Baths | Conduction | Superficial | Arthritis of small joints | Dilek et al., 2013 |
| Ultrasound Therapy | Conversion | 3–5 cm | Tendons, ligaments, joint capsules | Nascimento et al., 2022 |
| Shortwave Diathermy | Conversion | Deep tissues | Large muscle groups, chronic pain | Nascimento et al., 2022 |

(2008) showed that stretching preceded by ultrasound-assisted heating was a massively better intervention in terms of ROM. More recently, heat and manual therapy were demonstrated to accelerate the recovery of post-rotator cuff surgery patients (Kim et al., 2023).

### Post-acute Stages of Healing

The acute to subacute conversion in the process of conversion favors angiogenesis, fibroblast and collagen remodeling, which helps in functional recovery on the part of the heat. However, during the acute inflammatory phase, heat can exacerbate edema and tissue injury. The available guidelines emphasize the importance of taking care of time, keeping some heat to be used during the repair and remodeling steps (Vrindten et al., 2025).

### Pre-exercise Warm-up and Stretching

Heat increases the pliability of the muscles and joint viscosity, making them stretch and preventing injury. It is also commonly applied in sports rehabilitation and preemptive conditioning. A recent trial corroborated that pre-stretching of hamstrings with heat locally increased hamstring flexibility and sprinting in athletes (Davis et al., 2020).

## CONTRAINDICATIONS, PRECAUTIONS, AND PATIENT EDUCATION

Although heat therapy can be regarded as safe under most circumstances, its improper use can result in negative effects that include delayed healing, burns, or worsening of the underlying pathology. Therefore, a complete clinical examination is necessary before treatment.

### Absolute Contraindications

Heat should not be used in a number of conditions. It should not be applied in case of acute trauma or inflammation, loss of sensation of neuropathy, or spinal cord active hemorrhage, injury, or malignancy in the field of treatment. On the same note, patients who are severely arterially insufficient, vascular impaired, with thrombophlebitis and those with deep vein thrombosis must not receive any thermal intervention, open wounds, infected tissue, or cases where the patient has cognitive impairment preventing them from recognizing or reporting adverse effects are also an absolute contraindication.

### Precautions

Elsewhere, the heat treatment can be pragmatic; however, only with care and close observation. An example is the avoidance of heating the abdomen, pelvic, or lumbar area in pregnant patients. Patients with cardiovascular disease, unregulated hypertension, or peripheral vascular disease must be especially evaluated because of the changed circulatory issues. Extremities with edema, as well as implants made of metal (especially those treated with shortwave diathermy or ultrasound), require extra attention. Another significant factor is age since pediatric patients and elderly ones tend to have low thermoregulation, which predisposes them to burns or overheating (Lehmann et al., 1974; Vrindten et al., 2025).

### Patient Education

Patient education of anticipated sensations and safe use is an essential part of the treatment. They are to feel a slow warmth and relaxation of the muscles, and temporary erythema might occur, although it will disappear within a few minutes. In order to reduce the risk of harm, clinicians need to apply sufficient protective layers and restrict exposure time to approximately 15 to 20 min. in superficial modalities and prevent long-term or unsupervised application, such as sleeping with the heating pad. The patients are to be sensitised as well regarding the potential side effects, such as burns, skin irritation, or blisters. Its use in acute inflammations can increase swelling and heating of large body parts, which can lead to dizziness or hypotension (Nadler et al., 2002; Alghadir et al., 2019).

## EFFICACY AND INTEGRATION INTO PHYSICAL THERAPY

Evidence-based practice should guide the incorporation of heat therapy in rehabilitation as it integrates clinical expertise, research findings and patient preferences. An accumulating body of clinical research upholds the use of thermotherapy in supplementing musculoskeletal rehabilitation.

### Evidence-Based Efficacy

Heat treatment has always proved to be effective in many fields. It plays a significant part in the alleviation of pains, and research indicates that continuous heat wraps are an effective treatment of chronic low back pain and that paraffin therapy significantly lessens pain caused by arthritis (Nadler et al., 2002; Dilek et al., 2013). Compared to the control group, pre-stretching ultrasound enhances the flexibility of joints in adhesive capsulitis, whereas heat-assisted stretching increases the recovery of post-surgical shoulder rehabilitation (Dogru et al., 2008; Kim et al., 2023). There are also signs of an increase in hand functioning, with fluid therapy being demonstrated to increase grip strength and decrease stiffness in osteoarthritis and the post-operative period (Peng et al., 2020). In addition to these particular results, localized heating has also been associated with better quality of life through decreased muscle soreness, enhanced comfort, and, overall, enhanced daily functionality among patients with chronic musculoskeletal disorders (Petrofsky et al., 2015; Hill et al., 2024). Regardless of the accumulating evidence of the clinical utility of HT in treating MSK pain and the examples of meta-analyses, systematic reviews, consensus papers, and randomized multicenter clinical trials involving patients with low back and neck (Freiwald et al., 2021; Lubrano et al., 2023; Hotfiel et al., 2024), knee, and wrist pain (Freiwald et al., 2021) as studies indicate a positive response (Table 3).

### Integration into Treatment Plans

Its best use is as part of a complete rehabilitation program. It is used before exercise and increases muscle elasticity and minimizes the possibility of strain when straining. In combination with manual therapy, it also reduces the protein muscle guarding, allowing more profound mobilization and more productive joint methods. Thermotherapy has been used

**Table 3.** Evidence-based effects of heat therapy: supporting modality

| Outcome | Supporting Modality | References |
|---|---|---|
| **Pain Reduction** | Heat wraps, paraffin baths | Nadler et al., 2002; Dilek et al., 2013 |
| **Improved Mobility** | Ultrasound + stretching | Dogru et al., 2008 |
| **Grip Strength** | Fluidotherapy | Peng et al., 2020 |
| **Quality of Life** | Superficial moist heat | Petrofsky et al., 2015 |

in combination with strategies of chronic pain treatment with pharmacological treatments, cognitive behavioral therapy, and lifestyle alterations in multidisciplinary approaches to chronic pain. Besides, it can be an effective addition to such modalities as electrical stimulation, massage, hydrotherapy, and strengthening exercises and can maximize the short- and long-term rehabilitation outcomes (Vrindten et al., 2025).

### Research Gap

Despite its widespread use, heat therapy lacks comprehensive guidance in clinical practice, particularly regarding indications, contraindications, precautions, and routes of application. This educational gap has been identified among various healthcare professionals, including pharmacists (Hotfiel et al., 2024). Furthermore, there are few large-scale, multicenter randomized controlled trials investigating long-term outcomes, and standardized dosing protocols such as optimal temperature, duration, and frequency are largely absent. Research on heat therapy in neurological rehabilitation and systemic diseases remains limited, and there has been minimal exploration of digital or portable heat devices and tele-rehabilitation applications (Alghadir et al., 2019). These limitations underscore the need for more robust evidence and clearer clinical guidelines to optimize the safe and effective use of heat therapy.

Heat therapy continues to play a very critical role in the current rehabilitation with much of its low cost, ease of use, and varied therapeutic advantages. It not only helps reduce pain and stiffness but also helps increase joint mobility, muscular flexibility, and overall functional restoration. Besides those physiological implications, thermotherapy has some significant psychological impact; numerous patients feel relaxed, comfortable, and well-being during the course of the treatment, which subsequently increases their compliance with the rehabilitation programs. This dedication is essential because patients who take an active part in their treatment have more chances to attain long-term positive results.

Therapeutic effects are usually the greatest when heat is used sparingly, particularly when the subacute and chronic stages of musculoskeletal healing are involved, where tissues are most receptive to heat. In addition, thermotherapy is rarely used alone; its use offers the most advantages when combined with exercise, stretching or manual therapy. This kind of combination approach not only enhances physical outcomes, but it is also compatible with a comprehensive approach to care that is widely used with multidisciplinary approaches to pain management.

Despite centuries of the usage of heat therapy, which continues to be one of the most popular interventions, the clinical protocols still lack serious standardization gaps. Modern study emphasizes the need to establish specific parameters of safe and effective use, such as optimum temperatures, frequency of treatment, and length of session in different groups of patients. The research directions of the future need to focus on large randomized controlled trials with long-term follow-up to produce greater evidence on the lasting functional improvement and quality-of-life improvements. Also, wearable technologies and digital health platforms are being advanced, which present new possibilities of providing personalized, continuous, in-home-based thermotherapy, which could change the nature of patient-centered rehabilitation. Additional emerging applications are in neurological care, e.g., motor recovery following stroke and systemic disorders with poor metabolism, circulation, or pain regulation. When all of the mentioned points are combined, it points to the current evidence as showing that, given proper patient selection, individual treatment planning, and strict adherence to safety measures, heat therapy will remain a useful clinical tool. In addition to an increased rate of physical recovery, it is also associated with quality-of-life improvements, which further supports its status as a fundamental pillar of modern rehabilitation interventions, even years after its introduction.

## CONCLUSION

Thermotherapy still remains a safe, inexpensive and diverse instrument in musculoskeletal therapy. Such non-invasive procedures as hot packs and paraffin and invasive ones such as ultrasound, are performed by stimulating blood circulation, relieving muscle contractions, and softening connective tissues. These effects are useful in relieving pain and increasing functioning, especially when rehabilitation is used in conjunction with exercise or manual treatment in later stages of recovery. The use of this type of benefit should be accompanied by exercise or manual therapy, but the dosage and contraindications should be taken into account. Even the work in the future must focus on improving the standards of treatment and finding new solutions, such as wearable or home-based equipment, so that heat therapy will be one of the consistent elements of contemporary treatment.

## REFERENCES

Alghadir AH, S Anwer, B Sarkar et al., 2019. Effect of 6-week retro or forward walking program on pain, functional disability, quadriceps muscle strength, and performance in individuals with knee osteoarthritis: a randomized controlled trial (retro-walking trial). BMC Musculoskeletal Disorders 20:159.

Blyth FM, AM Briggs, CH Schneider et al., 2019. The global burden of musculoskeletal pain—where to from here? American Journal of Public Health 109:35–40.

Brunt VE and CT Minson, 2021. Heat therapy: mechanistic underpinnings and applications to cardiovascular health. Journal of Applied Physiology 130:1684–1704.

Charkoudian N, 2010. Mechanisms and modifiers of reflex-induced cutaneous vasodilation and vasoconstriction in humans. Journal of Applied Physiology 109:1221–28.

Clijsen R, R Stoop, E Hohenauer et al., 2022. Local heat applications as a treatment of physical and functional parameters in acute and chronic musculoskeletal disorders or pain. Archives of Physical Medicine and Rehabilitation 103:505–22.

Davis S, S Alabed, TJA Chico et al., 2020. Effect of sports massage on performance and recovery: a systematic review and meta-analysis. BMJ Open Sport and Exercise Medicine 6:000614.

Dilek BM, M Gözüm, S Gülbahar et al., 2013. Efficacy of paraffin bath therapy in hand osteoarthritis: a single-blinded randomized controlled trial. Archives of Physical Medicine and Rehabilitation 94:642–49.

Dogru HS, H Basaran, T Sarpel et al., 2008. Effectiveness of therapeutic ultrasound in adhesive capsulitis. Joint Bone Spine 75:445–50.

Freiwald J, E Paulino, G Solarino et al., 2021. A role for superficial heat therapy in the management of non-specific, mild-to-moderate low back pain in current clinical practice: a narrative review. Life 11:780.

Gianfaldoni S, U Wollina, MG Roccia et al., 2017. History of the baths and thermal medicine. Open Access Macedonian Journal of Medical Sciences 5:566.

Hill N, N Sharafi, A Clegg et al., 2024. The effectiveness and safety of heat/cold therapy in adults with lymphoedema: systematic review. Disability and Rehabilitation 46:2184–95.

Hotfiel T, M Blondett, J Freiwald et al., 2024. Importance of heat therapy in the treatment of pain in daily clinical practice. Journal of Bodywork and Movement Therapies 38:263–68.

Kim SJ, JH Choi, J Jung et al., 2023. Effectiveness of paraffin bath therapy for the symptoms and function of hand diseases: a systematic review and meta-analysis of randomized controlled trials. Journal of Hand Therapy 36:706–12.

Lehmann CG, CG Warren, SM Scham et al., 1974. Therapeutic heat and cold. Clinical Orthopaedics and Related Research 99:207–245.

Li Y, X Chen, X Ding et al., 2023. Exercise intervention for patients with chronic low back pain: a systematic review and network meta-analysis. Frontiers in Public Health 11:1155225.

Lubrano E, I Grattagliano, LS de Medeiros et al., 2023. An international multidisciplinary Delphi-based consensus on heat therapy in musculoskeletal pain. Pain and Therapy 12:93–110.

Nadler DA, DA Hengehold, RW Weingand et al., 2002. Continuous low-level heat wrap therapy provides more efficacy than ibuprofen and acetaminophen for acute low back pain.

dos Reis do Nascimento MR, GC Raab Ferreira, LH Silva Goncalves et al., 2022. Immediate analgesic effect of two modes of short-wave diathermy application in chronic low back pain: study protocol for a randomized controlled trial. Pain Management 12:131-39.

Peng L, M Chi, LM Lin et al., 2020. The effectiveness of far-infrared irradiation on foot skin surface temperature and heart rate variability in healthy adults over 50 years of age: a randomized study. Medicine 99:23366.

Petrofsky J, F Al-Dabbak, MS Laymon et al., 2015. Cold vs heat after exercise—is there a clear winner for muscle soreness? Journal of Strength and Conditioning Research 29:3245–52.

Rossi R, 2024. Heat therapy for different knee diseases: expert opinion. Frontiers in Rehabilitation Sciences 5:1390416.

Rutkove SB, 2001. Effects of temperature on neuromuscular electrophysiology. Muscle and Nerve 24:867–882.

Ventriglia A, G Magni, G Panico et al., 2023. Musculoskeletal pain management and thermotherapy: an exploratory analysis of Italian physicians' attitudes, beliefs, and prescribing habits. Journal of Pain Research 16:1547–57.

Vrindten JL, JL Mazzocca, EG Matzkin et al., 2025. Thermal modalities including hot baths and cold plunges play a unique role in injury prevention and recovery. Arthroscopy, Sports Medicine and Rehabilitation 7:101143.